

STELLAR CONVECTION IN LATE-TYPE GIANTS AND SUPERGIANTS

Alex Brown (CASA, University of Colorado)

1. Convection in the Sun and stars
2. Stellar Dynamos
3. Solar and stellar activity
4. Stellar surface structure
5. Potential for interferometric observations

"CLASSICAL" CONVECTION

Schwarzschild criterion for convective instability

$$\left| \frac{dT}{dr} \right| > \left| \left(\frac{dT}{dr} \right) \right|_{\text{adiabatic}}$$

BOHM-
VITENSE
(1958, Zs. f. Ap
46, 108)

Mixing length $\alpha = \frac{L_{ML}}{H_p}$ — mixing length $\alpha \approx 1.6$
 H_p — pressure scale height

Convective flux density


$$F_c \approx \frac{5}{2} \rho(r) \tilde{v}^3(r)$$

$$\equiv \sigma T_{\text{eff}}^4 \frac{R^2}{r^2}$$

if convection is
only transport mech.

Radiative flux density

$$F_R = - \frac{16 \sigma T^3}{3 \kappa_R \rho} \frac{dT}{dr}$$



SOLAR CONVECTION ZONE - IN CONTEXT

$R_{\odot} \sim 700 \text{ Mm}$

Core $\sim 100 \text{ Mm}$

50% of mass $R < 175 \text{ Mm}$

Radiative Zone $\sim 500 \text{ Mm}$

$[R = 0.71 - 1.0 R_{\odot}]$ Convection Zone $500 \rightarrow 700 \text{ Mm}$ $\frac{2}{3}$ volume
 $\frac{2}{6}$ of mass

Photosphere $\approx \text{few} \times 10^2 \text{ m}$ thick

Chromosphere $\sim 10 \text{ Mm}$ thick

Corona - many scales $(0.1 R_{\odot} - 1 R_{\odot})$

Solar Wind $\gg 1 \text{ AU}$.

CONVECTION IN OTHER STARS

Models at use mixing length theory

→ stellar structure models

→ stellar evolution models

"Cool stars" \equiv those with convection zones just below photosphere

Thin convection zone - late A / early F stars

Fully convective - mid M stars

KM Giants / Supergiants - deepening CZ
- convective mixing


[Evolution models need convective overshoot]

Solar Supergranulation and Surface Flows

High resolution SOHO - MDI magnetogram overlaid with lines of convergence of the photospheric horizontal flows.

- Measured flow shown as colored arrows; red = downflow, blue = upflow.
- Green dots mark the convergence points.
- Magnetic field; light grey for positive fields, for negative fields.

23 Feb. 1996, 16:44 to 21:03 UT



NUMERICAL 3-D MHD SIMULATIONS OF SOLAR CONVECTION

Work of e.g. Nordlund, Stein, Dravins,
Toomre, Brummel

Labinitio 3D MHD - field equations, radiative transfer, atomic physics

Heroic effort - new insights into how
convection operates in outer few
pressure scale heights of Sun

They find - upward and downward moving
elements do NOT mix.

- circulation driven by cooling at surface
NOT by buoyant bubbles.

- large upflow areas feed strong
downdrafts.


Flow scales: ~ 10 min at top of CZ, ~ 1 month at bottom of CZ

Excellent match to temperature and velocity fields,
photospheric absorption line profiles. Also fit p modes

Full problem still too hard


Density range 10^6 , pressure range 10^9

surface brightness; 3-dimensional simulations




253 x 253 x 163 125 x 125 x 105

as above, simulated 40 cm telescope & atmosphere



253 x 253 x 163 125 x 125 x 105

observations, Swedish 40 cm @ La Palma



Solar Convection


- **Global / Convection Zone**

- Inner 3-4 orders of magnitude in density
 - 95% of CZ depth
- Scales of the order of CZ depth
- Turn-over times ~ months
- Polytropic stratification
 - index $\sim 3/2$, nearly ideal




- **Surface layers**


- Outer 4-6 orders of magnitude in density
 - 5% of CZ depth ~ 10 Mm
- Scales from several $\times 10$ Mm to below 1 Mm
- Turn-over times from days to minutes
- Stratification: ~exponential
 - polytropic with large index



Fortunately, life is more interesting



G Band - Swedish Vacuum Solar Telescope, La Palma, Spain.




← 132 Mm →

G-Band

AR 9037, 14 June 2000, 10:00 UT

Swedish Vacuum Solar Telescope, La Palma, Spain

L. Strous, D. Torgerson, Stanford Lockheed Institute for Solar Research



H α Line Center

AR 9037, 14 June 2000, 10:00 UT

Swedish Vacuum Solar Telescope, La Palma, Spain

L. Stenou, D. Torgerson, Stanford Lockheed Institute for Solar Research

Stellar Dynamos

DIFFERENTIAL ROTATION + TURBULENT CONVECTION

⇒ MAGNETIC FIELD GENERATION

For Sun - $\alpha\Omega$ dynamo - operating in shear zone
at bottom of convection zone

Magnetic flux stored in "under shoot layer"

Convection inhibits magnetic buoyancy.

Strongest field at bottom of CZ - $\text{few} \times 10 \text{ KG}$

"Turbulent" dynamos also operate in Sun and other stars
[α^2]


Details of dynamo theory largely untested observationally

For Sun theories try to match

- 22yr solar cycle + polarity reversal
- spot migration - "Butterfly Diagram"
- differential rotation
- behavior of emerging field into photosphere

Other stars can show RADICALLY different activity patterns

Manifestations and Effects of Stellar Magnetic Activity



- **Solar luminosity shows cyclic changes**
 - induced climate changes on Earth, such as the 17th-Century Little Ice Age during the solar Maunder minimum
- **In solar/stellar atmospheres:**
 - magnetic regions & star spots;
 - very hot outer atmospheres;
 - explosive flares & high-energy particles and radiation;
 - stellar wind & coronal mass ejections

Stellar Oscillations


Source: *Lecture Notes on Stellar Oscillations*, Jorgen Christensen-Dalsgaard
(Aarhus University, Denmark) – <http://www.obs.aau.dk/jcd/oscilnotes/comb-h.tex.ps.gz>

MDI Imaging of the Sun

- Single Dopplergram from Michelson Doppler Imager (MDI) on SOHO.
- MDI measures the photospheric radial velocity.
- Dominant effect in single Dopplergram is solar rotation.

Single Dopplergram

(30-MAR-96 19:54:00)




MDI Imaging of the Sun


- Subtracting an average Dopplergram (summed over 45 minutes) from a single velocity image reveals the surface motions associated with sound waves traveling through the Sun's interior.
- The smallscale light and dark regions represent the up and down motions near the solar surface.
- The pattern falls off towards the limb because the acoustic waves are primarily radial.



Single Dopplergram Minus 45 Images Average

(30-MAR-96 19:54:00)



MDI Medium- l Power Spectrum








Figure 5.6. Cyclic frequencies $\nu = \omega/2\pi$, as functions of degree l , computed for a normal solar model. Selected values of the radial order n have been indicated.





Figure 2.1. Contour plots of the real part of spherical harmonics Y_l^m [cf. equation (2.1); for simplicity the phase factor $(-1)^m$ has been suppressed]. Positive contours are indicated by continuous lines and negative contours by dashed lines. The $\theta = 0$ axis has been inclined by 45° towards the viewer, and is indicated by the star. The equator is shown by “++++”. The following cases are illustrated: a) $l = 1, m = 0$; b) $l = 1, m = 1$; c) $l = 2, m = 0$; d) $l = 2, m = 1$; e) $l = 2, m = 2$; f) $l = 3, m = 0$; g) $l = 3, m = 1$; h) $l = 3, m = 2$; i) $l = 3, m = 3$; j) $l = 5, m = 5$; k) $l = 10, m = 5$; l) $l = 10, m = 10$.


Solar Internal Rotation

- Inferred rotation rate as a function of depth and latitude derived from SOHO helioseismology.
- Convection zone rotates uniformly along a radius with all depths showing the surface differential rotation.
- Below the convection zone is the shear layer of shear connecting to the rigidly rotating radiative core.



Solar Internal Temperature Distribution


- Red layers - sound travels faster than predicted by the theories, implying that the temperature is higher than expected.
- Blue layers - sound speed (and temperature) is lower than expected.
- Higher temperatures than expected seen at the transition between the convection zone and the radiative core. This shear zone, between the faster-turning outer region and the slower interior, is directly related to generating the magnetic dynamo.
- The core is cooler than predicted by $\sim 0.1\%$. Although the discrepancy may seem small, it implies that the thermonuclear energy generation could be underperforming at present.



[Credit: SOHO (ESA & NASA), MDI/SOI and VIRGO data imaged by A. Kosovichev, Stanford University]


STARSPOTS - Doppler Imaging

Most information on starspot distributions on active stars has come from Doppler Imaging studies. As a starspot moves across the stellar disk it appears as a distinct feature in the optical absorption line spectrum.



STARSPOTS - Doppler Imaging

Doppler images of UZ Lib (K0 III, rotational period 4.7 days) constructed by Strassmeier (1996, A&A 314, 558) based on data from 1994 March. These images show many features typical of active stars: strong photospheric temperature contrasts with cool spots spread over the stellar surface, large spot filling factors, the presence of polar spots.



SURFACE STRUCTURE ON RED SUPERGIANTS



- Red giants and supergiants show only weak magnetic activity.
- Their atmospheres are dominated by wind outflow.
- Do not expect large starspots as on the Sun or active stars.
- Supergiants may show giant convective cells in their photospheres - suggested by Schwarzschild (1975 ApJ 195, 137).

SURFACE STRUCTURE - HST UV Imaging

- HST imaging using FOC and other cameras shows asymmetric light distribution in Mg II 2800Å emission (Uitenbroek et al. .
- Size of disk is $\sim 2 R_*$.
- This emission is being formed in base of wind.

Mg II emission to 270 m.
from long slit spect
(GHRs)

1995
March




$d \approx 131 \text{ pc}$
 $R_* \sim 800 R_\odot$

SURFACE STRUCTURE - Optical/IR Interferometry

- Optical interferometry of a range of red supergiants frequently detects asymmetric light distributions - "hot spots".
- "Hot spots" are seen to be time variable and sometimes are not present at all.
- As an example, below, are reconstructed images of Betelgeuse 7000Å and 9050Å (Young et al. 2000 MN 315, 635).
- Presently, it is very unclear whether these correspond in any way to the Schwarzschild convective cells.


1997
05/NOV



SURFACE STRUCTURE - Aperture Masking

- Non-redundant aperture masking using large telescopes provides similar images.
- Another example of Betelgeuse from Young et al. (2000).


1997
NOV



SURFACE STRUCTURE - Radio Interferometry

- Radio images of supergiants show nonuniform emission.
- Lim et al. found an asymmetric disk at 7 mm for Betelgeuse.
- This emission is extended and formed in regions of similar size to the UV emission lines.


7mm Radio Image of
Betelgeuse's Atmosphere





Courtesy of J. Lim, C Carilli, S. M. White,
A. J. Beasley, & R. G. Marson



(1998, *Nature* 392, 575)

Visibility (mJy)












Figure 5.4. Propagation of acoustic waves, corresponding to modes with $l = 30$, $\nu = 3$ mHz (deeply penetrating rays) and $l = 100$, $\nu = 3$ mHz (shallowly penetrating rays). The lines orthogonal to the former path of propagation illustrate the wave fronts.

Stellar angular diameters:

Largest M supergiants $\sim 40-60$ mas

Largest K giants ~ 20 mas

Active binary primaries $\sim 10-20$ mas

Nearby dwarf stars \sim few mas